arXiv:1101.4444v2 [math.DG] 7 Jul 2011

A RESTRICTION FOR SINGULARITIES ON COLLAPSING ORBIFOLDS

YU DING

Abstract. Every point p in an orbifold X has a neighborhood that is homeomorphic to $G_p \backslash B_r(0)$, where G_p is a finite group acting on $B_r(0) \subset \mathbb{R}^n$, so that $G_p(0) = 0$. Assume X is a Riemannian orbifold with isolated singularities that is collapsing, that is, X admits a sequence of metrics g_i with uniformly bounded curvature, so that for any $x \in X$, the volume of $B_1(x)$, with respect to the metric g_i, goes to 0 as $i \to \infty$. For such X, we prove that $|G_p| \leq (2\pi/0.47)^{n(n-1)}$ for all singularities $p \in X$.

1. Introduction

An n dimensional Riemannian *orbifold*, X, is a metric space so that the following is true: for any $x \in X$, there exists $r = r(x) > 0$ and a Riemannian metric $\tilde{g}_x$ on $B_{2r}(0) \subset \mathbb{R}^n$, a finite group G_x (the isotropy group) acting on $(B_r(0), \tilde{g}_x)$ by isometries, so that $G_x(0) = 0$, and there is an isometry $\iota_x : B_r(x) \to G_x \backslash B_r(0)$ with $\iota_x(x) = 0$. See [Th]. We call $x \in X$ a *regular point* if $|G_x| = 1$; otherwise, x is a *singular point*. We say the curvature of X satisfies

$$|K_X| \leq \kappa^2, \tag{1.1}$$

if the sectional curvature K of every $(B_r(0), \tilde{g}_x)$ above satisfies $|K| \leq \kappa^2$. We say X is *collapsing*, if X admits a sequence of metrics, g_i, with uniformly bounded curvature, so that for any $x \in X$,

$$\lim_{i \to \infty} \mathrm{Vol}_{g_i}(B_1(x)) = 0. \tag{1.2}$$

As an example, consider the standard $\mathbb{Z}_m = \mathbb{Z}/m\mathbb{Z}$ action on the sphere S^2:



$X = \mathbb{Z}_m \backslash S^2$

The quotient orbifold $X = \mathbb{Z}_m \backslash S^2$ will be arbitrarily collapsed when $m \to \infty$. However, for any *fixed* m, (i.e. each of the two singularities there is a neighborhood that is isometric to $\mathbb{Z}_m \backslash \mathbb{R}^2$, where $\mathbb{R}^2$ is equipped with some $\mathbb{Z}_m$ invariant metric). $X = \mathbb{Z}_m \backslash S^2$ can be collapsed only to a certain degree, it does not support a sequence of collapsing metrics; in fact, otherwise by pull back we get a sequence of collapsing metrics on S^2, a contradiction to the Gauss-Bonnet theorem.

Date: July 8, 2011.

On the other hand, consider the double of a 2-dimensional rectangle. Clearly it admits a flat metric, thus we obtain a sequence of collapsing metrics by rescale. Notice, in this example, for each of the four singularities, the isotropy group G_x has order 2, a quite small number.

The main result of this paper is

Theorem 1.3. *Assume X is a compact, collapsing orbifold, $p \in X$ is an isolated singularity. Then $|G_p| \leq (2\pi/0.47)^{n(n-1)}$.*

If X has an isolated singularity p, then the dimension of X must be even, and $G_p \subset SO(n)$. The bound $|G_p| \leq (2\pi/0.47)^{n(n-1)}$ has its root in the Bieberbach theorem of crystallographic groups and Gromov's almost flat manifold theorem; however, it fails when we drop the requirement that x is an *isolated* singularity. For example, we can take *any* orbifold X' and let $X = X \times S^1$; by shrinking the S^1 factor, we see X is collapsing while there is no restriction on singularities of X'.

Clearly, Theorem 1.3 is a corollary of the following:

Theorem 1.4. *For any $L > 0$, there is $\epsilon = \epsilon(n, L)$ so that if X is an orbifold with all singularities $q \in X$ satisfying $|G_q| < L$, $\mathrm{Vol}(B_1(q)) < \epsilon$, then $|G_p| \leq (2\pi/0.47)^{n(n-1)}$ for any isolated singularity $p \in X$.*

A *nilmanifold*, $\Gamma\backslash N$, is the quotient of the (left) action of a discrete, uniform subgroup $\Gamma \subset N$, on a simply connected nilpotent Lie group N. Left invariant vector fields (LIVFs) can be defined on $\Gamma\backslash N$. An affine diffeomorphism of $\Gamma\backslash N$ is a diffeomorphism that maps any local LIVF to some local LIVF. In general, a right invariant vector field (RIVF) cannot be defined globally in $\Gamma\backslash N$, unless this vector field is in the center of the Lie algebra of N. However, the right invariant vector fields, not the left invariant ones, are Killing fields of left invariant metrics on $\Gamma\backslash N$. An *infranil orbifold* is the quotient of a nilmanifold by the action of a finite group H of affine diffeomorphisms. If the action H is free, we get an *infranil manifold.*

In our previous work, [D], we generalized the Cheeger-Fukaya-Gromov nilpotent Killing structure, [CFG], and the Cheeger-Gromov F-structure, [CG1], [CG2], to collapsing orbifolds. In particular, sufficiently collapsed X can be decomposed into a union of *orbits.* Each orbit $\mathscr{O}_p$ is the orbit of the action of a sheaf $\mathfrak{n}$ of nilpotent Lie algebras, which comes from local RIVFs on a nilmanifold fibration in the frame bundle FX. Therefore every $\mathscr{O}_p$ is an infranil orbifold. The proof of Theorem 1.4 is based on the relation between singularities on X and singularities within an orbit $\mathscr{O}_p$ in X, as well as the nilmanifold fibration on FX.

X is called *almost flat*, if

$$\sup |K_X|^{1/2} \cdot \mathrm{Diam}\, X \leq \delta_n, \tag{1.5}$$

here $\mathrm{Diam}\, X$ is the diameter of X, δ_n is a small constant that depends only on n. In [Gr], Gromov proved that an almost flat *manifold* M has a finite, normal covering space $\tilde{M} = \Gamma\backslash N$ that is a nilmanifold. Subsequently, Ruh, [Ru], proved that M is diffeomorphic to $\Lambda\backslash N$, where $\Lambda \supset \Gamma$ is a discrete subgroup in the affine transformation group of N. In [Gh2], Ghanaat generalized this to an almost flat orbifold X, under the assumption that X is *good* in the sense of Thurston [Th], i.e. X is the *global* quotient of a simply connected manifold M. There are examples of orbifolds that are not good, see [Th]. In fact, without much effort, one can remove the assumption that X is good:

Theorem 1.6. *If X is an almost flat orbifold, then X is an infranil orbifold.*

Precisely, there is a nilmanifold $\tilde{X} = \Gamma \backslash N$, a finite group H acting on $\tilde{X}$ by affine diffeomorphism, so that X is diffeomorphic to $H \backslash \tilde{X}$. The order of H is bounded by $c_n \leq (2\pi/0.47)^{n(n-1)/2}$. Moreover, there is a sequence of metrics g_j so that $\mathrm{Diam}(X, g_j) \to 0$.

The proof of Theorem 1.4 does not depend on the above result, so we only give a sketch of its proof in the Appendix. On the other hand, this implies Theorem 1.4 for almost flat orbifolds immediately, even without the assumption that the singularities are isolated.

2. Proof of Theorem 1.4

If X is an infranil orbifold, then it is easy to obtain the bound in Theorem 1.3. Since the proof contains some ideas for the general case, we give full details.

Lemma 2.1. *Assume X is an infranil orbifold. Then $|G_x| \leq (2\pi/0.47)^{n(n-1)/2}$.*

Proof. Assume $X = \Lambda \backslash N$, where N is a simply connected nilpotent Lie group, Λ is a discrete group of affine diffeomorphisms on N so that $X = \Lambda \backslash N$ is compact. If N is abelian, then X is a flat orbifold, Λ is a discrete group of isometries on $N = \mathbb{R}^n$ that acts properly discontinuously. So the conclusion follows from (the proof of) Bieberbach's theorem on crystallographic groups. In fact, it is well known that the maximal rotational angle of any $\lambda \in \Lambda$ is either 0 or at least 1/2. Thus the bound comes from a standard packing argument; notice $n(n-1)/2 = \dim SO(n)$ and the bi-invariant metric on $SO(n)$ has positive curvature.

We prove the general case by induction on dimension of X. Remember that Λ contains a normal subgroup Γ of finite index, so that Γ is a uniform, discrete subgroup of N. X the the quotient of the Λ/Γ action on the nilmanifold $\tilde{X} = \Gamma \backslash N$. Clearly G_x embeds in Λ/Γ, that is, $G_x = \{\bar{\lambda} \in \Lambda/\Gamma \mid \bar{\lambda}\tilde{x} = \tilde{x}\}$; here we choose a point $\tilde{x}$ in $\tilde{X} = \Gamma \backslash N$ that projects to $x \in \Lambda \backslash N$.

Let C be the center of N, then C is connected, of positive dimension. Since any $\lambda \in \Lambda$ is affine diffeomorphism, λ moves a C-coset in N to a C-coset. Therefore Λ/Γ acts on the nilmanifold $\tilde{X}^* = (\Gamma/(\Gamma \cap C)) \backslash (N/C)$, the quotient X^* is an infranil orbifold of lower dimension. Let $\pi : \tilde{X} \to \tilde{X}^*$ be the projection, assume $\pi(\tilde{x}) = \tilde{x}^*$. Thus we have a homomorphism

$$h : G_x \to G_{x^*}. \tag{2.2}$$

$\tilde{X}$ is a torus bundle over $\tilde{X}^*$, the fiber is $T = (\Gamma \cap C)\backslash C$. Assume $\bar{\lambda} \in \Lambda/\Gamma$ is in Ker h, the kernel of h, then $\bar{\lambda}$ fixes every T fiber in $\tilde{X}$. If, in addition, $\bar{\lambda}$ fixes *every point* in the T fiber passing through $\tilde{x}$, we claim $\bar{\lambda}$ must be identity. In fact, on N we have $\lambda(z) = a \cdot A(z)$, here $a \in N$ and A is a Lie group automorphism of N; if $\bar{\lambda}$ fixes every point in *one* T fiber, then A is identity on the center $C \subset N$. This implies that $\bar{\lambda}$ is a *translation* on every T fiber. Since $\bar{\lambda}$ is of finite order and fixes every point in *one* T-fiber, $\bar{\lambda}$ must be identity. Therefore any element $\bar{\lambda} \in$ Ker h is decided by its restriction on the T fiber passing through x; so Ker h is isomorphic to a finite group of affine diffeomorphisms on T that fixes $\tilde{x} \in T$, thus |Ker h| can be bounded by Bieberbach's theorem. Since

$$|G_x| \leq |G_{x^*}| \cdot |\text{Ker } h|, \tag{2.3}$$

the conclusion follows by induction. □

In [D], the existence of nilpotent Killing structure of Cheeger-Fukaya-Gromov, [CFG], is generalized to sufficiently collapsed orbifolds. We briefly review this construction.

As in the manifold case, one can define the *frame bundle* FX of an orbifold X. If $B_r(x) \subset X$ is isometric to $G_x\backslash B_r(0)$, where G_x is a finite group acting on $B_r(0) \subset \mathbb{R}^n$, then locally FX is $G_x\backslash FB(0,r)$, here $FB(0,r)$ is the orthonormal frame bundle over $B_r(0)$, and G_x acts on $FB(0,r)$ by differential, i.e. $\tau \in G_x$ moves a frame u to $\tau_* u$. Therefore FX is a manifold; strictly speaking, FX is not a fiber bundle. Let $\pi : FX \to X$ be the projection.

Moreover, there is a natural $SO(n)$ action on FX; on the frames over regular points, this $SO(n)$ action is the same one as in the manifold case; however, at (the frames over) singular points, this action is *not free.* As in the work of Fukaya, [F2], see also [D], any Gromov-Hausdorff limit Y of a collapsing sequence FX_i is a manifold. Following [CFG], for sufficiently collapsing orbifolds, *locally* we have an $SO(n)$-equivariant fibration

$$Z \longrightarrow FX \xrightarrow{f} Y, \tag{2.4}$$

here the fiber Z is a nilmanifold, Y is a smooth manifold with controlled geometry.

As in [CFG], we can put a *canonical* affine structure on the Z fibers, i.e. a canonical way to construct a diffeomorphism from a fiber Z to the nilmanifold $\Gamma\backslash N$. In particular, there is a sheaf $\mathfrak{n}$, of a nilpotent Lie algebra of vector fields on FX. Sections of $\mathfrak{n}$ are local right invariant vector fields on the nilmanifold fibers Z. By integrating $\mathfrak{n}$, we get a *local* action of a simply connected nilpotent Lie group, $\mathfrak{N}$, on FX. Therefore we also call a Z fiber an *orbit*, we can write $Z = \tilde{\mathscr{O}}$.

The fibration $f : FX \to Y$ is $SO(n)$-equivariant, so any $Q \in SO(n)$ moves a Z fiber to a (perhaps another) Z fiber by affine diffeomorphism. Moreover, the $SO(n)$ action on $\mathfrak{n}$ is *locally trivial*, that is, if $A \in \mathfrak{so}(n)$ is sufficiently small, then $e^A \in SO(n)$ moves a section, $\mathfrak{n}(U)$, of $\mathfrak{n}$ on any open set $U \subset FX$, to itself (over $U \cap Ue^A$). See [CFG] Proposition 4.3. In particular, the sheaf $\mathfrak{n}$ induces a sheaf, which we also denote by $\mathfrak{n}$, on the orbifold X away from the singular points. An orbit $\tilde{\mathscr{O}}_{\tilde{q}} = Z$ on X projects down to an *orbit* $\mathscr{O}_q$ on X.

Assume $\tilde{q} \in FX$ is any frame over $q \in X$. Let

$$I(q) = \{Q \in SO(n) \,|\, Z_{\tilde{q}}Q = Z_{\tilde{q}}\} \tag{2.5}$$

be the the isotropy group of an orbit $Z_{\tilde{q}} = \mathscr{O}_{\tilde{q}} \subset FX$. We will simply write $I(q)$ by I. Let I_0 be the identity component of I. It can be shown that restricted on $Z = \tilde{\mathscr{O}}_{\tilde{q}}$, the action of I_0 is identical to the action of a torus, and the Lie algebra of this torus, I_0, is in the center of $\mathfrak{n}$. See [CFG], and [D] for more details. Consider the nilmanifold

$$\check{\mathscr{O}}_{\tilde{q}} = \tilde{\mathscr{O}}_{\tilde{q}}/I_0. \tag{2.6}$$

Therefore $Z_{\tilde{q}} = \tilde{\mathscr{O}}_{\tilde{q}}$ is a torus bundle over $\check{\mathscr{O}}_{\tilde{q}}$. Notice, on $Z = \tilde{\mathscr{O}}_{\tilde{q}}$, I moves I_0 fibers to I_0 fibers, thus the orbit $\mathscr{O}_q$ is the quotient of $\check{\mathscr{O}}_{\tilde{q}}$ by the action of the finite group I/I_0. Therefore $\mathscr{O}_q$ is an infranil orbifold. In particular, the singularities within $\mathscr{O}_q$ satisfies the bound in Lemma 2.1.

It is important to remark that the above structure is not trivial:

Lemma 2.7. *Let L be any integer. Then there is $\epsilon = \epsilon(n, L)$, so that if X is an orbifold with $|G_x| \leq L$, $\operatorname{Vol}(B_1(x)) \leq \epsilon$ for all $x \in X$, then every $\mathfrak{n}$-orbit $\mathscr{O}$ on X is of positive dimension.*

Proof. (sketch) For any unit vector $A \in \mathfrak{so}(n)$, the bound in $|G_x|$ implies that e^{tA} does not have fixed point in $\tilde{\mathscr{O}}_{\tilde{q}}$ unless $t = 0$ or $|t| > cL^{-1}$. However, for sufficiently collapsed orbifolds, there is a vector B in the center of $\mathfrak{n}$ so that B generates a closed loop in $\tilde{\mathscr{O}}_{\tilde{q}}$ that is shorter than cL^{-1}, therefore B cannot be in the Lie algebra of I_0, which is in both $\mathfrak{so}(n)$ and the center of $\mathfrak{n}$. Thus the orbit $\tilde{\mathscr{O}}_{\tilde{q}}$ is not contained in a single $SO(n)$ orbit in FX, so $\mathscr{O}_q$ is of positive dimension in $X = FX/SO(n)$. See [D] for more details. ☐

Proof of Theorem 1.4. Assume $p \in X$ is an isolated singular point, $\tilde{p} \in \pi^{-1}(p)$ is in FX. $Z_{\tilde{p}} = \tilde{\mathscr{O}}_{\tilde{p}}$ is the fiber that projects to $\mathscr{O}_p$. Let $I(p)$, I_0, $\check{\mathscr{O}}_{\tilde{p}}$ be as above. Let

$$K_p = \{Q \in SO(n) \,|\, \tilde{p}Q = \tilde{p}\}. \tag{2.8}$$

Thus K_p is a subgroup of I, and $|K_p| = |G_p|$. Let

$$K_p^- = \{Q \in K_p \,|\, (\tilde{p}'I_0)Q = (\tilde{p}'I_0) \quad \text{for all } \tilde{p}' \in Z_{\tilde{p}}\}. \tag{2.9}$$

Thus K_p^- is a normal subgroup of K_p.

Lemma 2.10. *If $Q \in K_p$ fixes every point in $Z_{\tilde{p}}$, then Q is the identity in $SO(n)$.*

Proof. Potentially Q may fix every point in $Z_{\tilde{p}}$ while moving some points of FX that are outside $Z_{\tilde{p}}$. We will rule out this possibility.

By assumption, p is an isolated singularity. For any Q that is not identity, the connected component of the fixed point set of Q that passes through $\tilde{p}$ must project to p under $\pi : FX \to X$, because away from $\pi^{-1}(p)$ the $SO(n)$ action is free. Therefore

$\pi(Z_{\tilde{p}}) = p$ is a single point in X, this contradicts the fact that the $\mathfrak{n}$-orbits on X are of positive dimension; see Lemma 2.7. □

In particular, we have a *faithful* representation of K_p in the affine group of $Z_{\tilde{p}}$, i.e. we can identify K_p with the *restricted action* of the group K_p *on* $Z_{\tilde{p}}$.

Take any $Q \in K_p^-$ that is not identity in $SO(n)$. By definition Q fixes $\tilde{p}$. If Q fixes *every* point in $\tilde{p}I_0$, as in Lemma 2.1, Q is a translation on *every* I_0 fiber; because Q is of finite order and Q moves every I_0 fiber to itself, Q necessarily fixes every point in Z; thus Q is identity. So Q rotates the tangent plane of $\tilde{p}I_0$ at $\tilde{p}$. Therefore K_p^- is isomorphic to a finite group of affine diffeomorphisms on the torus $\tilde{p}I_0$. By the Bieberbach theorem,

$$|K_p^-| \leq (2\pi/0.47)^{k(k-1)/2}, \qquad k = \dim I_0. \tag{2.11}$$

Recall that Bieberbach's theorem implies that all finite subgroups of $SL(n, \mathbb{Z})$ have a uniform upper bound in order.

We have

$$\mathscr{O}_p = H \backslash (Z_{\tilde{p}}/I_0), \tag{2.12}$$

where $H = I/I_0$ is a finite group. Let H_p be the subgroup of H that fixes p. Now we get an *embedding*

$$K_p/K_p^- \subset H_p. \tag{2.13}$$

By Lemma 2.1,

$$|H_p| \leq (2\pi/0.47)^{i(i-1)/2}, \qquad i = \dim \mathscr{O}_p. \tag{2.14}$$

Thus

$$|G_p| = |K_p| = |K_p/K_p^-| \cdot |K_p^-| \leq (2\pi/0.47)^{n(n-1)}. \tag{2.15}$$

□

APPENDIX: ORBIFOLD VERSION OF GROMOV'S THEOREM

Assume M is an almost flat manifold. The proof of Gromov's almost flat manifold theorem has two natural steps. First, by a delicate calculus of geodesic loops, one gets a finite, normal covering $\hat{M}$ of M; see [Gr], [BK] Chapters 1-3. Next, one uses the deformation technique of Ruh [Ru] and Ghanaat [Gh] to show that $\hat{M}$ is a nilmanifold. We follow closely this approach. See also [Ro].

Rescale X so that $\operatorname{Diam} X = 1$, therefore the curvature of X is bounded by δ_n^2. Let $p \in X$ be a regular point, and v be any unit tangent vector at p. The geodesic γ so that $\gamma(0) = p$ and $\gamma'(0) = v$ is defined at least on a short interval $[0, \delta)$. One can extend it to be a geodesic $\gamma : [0, \infty) \to X$. In fact, the extension exists as long as $\gamma(t)$ is a regular point; if $\gamma(t) = x$ is singular, recall there is a neighborhood U_x of x, an Euclidean ball V_x with metric $\tilde{g}$ so that $U_x = G_x \backslash V_x$; we can lift $\gamma|_{(t-\epsilon, t]}$ to a geodesic on V_x, extend it beyond t on V_x and then project to $U_x \subset X$. Therefore, we get a map

$$\exp : \mathbb{R}^n = T_p \longrightarrow X, \tag{2.16}$$

so that $\exp(sv) = \gamma(s)$. This map is called the *develop map*; see [Th], [CT]. It plays the role of exponential map. Let

$$\rho < r_{\max} = \pi/\sqrt{\delta_n}. \tag{2.17}$$

By standard comparison theorem, exp is *nonsingular* on $B_\rho(0)$. When $x = \exp(tv)$ is a singular point this means, there is a neighborhood $\hat{V}$ of $tv \in \mathbb{R}^n$ so that the following local lift $\hat{\exp}$ is nonsingular (it is always smooth):

$$\hat{V} \xrightarrow{\hat{\exp}} V_x \xrightarrow{\mathrm{pj}_x} U_x \subset X, \qquad \exp|_{\hat{V}} = \mathrm{pj}_x \circ \hat{\exp}. \tag{2.18}$$

In particular, the Riemannian metric g on X pulls back to be a smooth metric $\tilde{g}$ on $B_\rho(0) \subset \mathbb{R}^n$. In the following, we will assume that $B_\rho(0) \subset T_p$, write T_p for simplicity, is equipped with the metric $\tilde{g}$.

We use the convention that for two loops α, β in X, the product $\alpha \cdot \beta$ is the loop that goes along β first, then α. Define

$$\pi_\rho = \{ \text{ geodesic loops } \gamma \text{ at } p \in X \text{ with length } |\gamma| \le \rho\}. \tag{2.19}$$

Choose ρ so that $2\,\mathrm{Diam}(X) \le 2 \ll \rho \ll r_{\max}$. Assume α, β are two loops at p, $|\alpha| + |\beta| < r_{\max}$. Then we can lift $\alpha \cdot \beta$ to a piecewise geodesic on T_p; such a path is homotopic (rel. end points) to a unique minimal geodesic $\tilde{\gamma}$ on T_p. Their *Gromov product*, $\alpha * \beta$, is defined to be $\exp(\gamma)$. This kind of homotopy on X, which can be lifted to T_p through exp, is called *Gromov homotopy.* In general $\alpha * \beta$ is a geodesic loop in X; but *not* the minimal geodesic in the homotopy class of $\alpha \cdot \beta$ - examples are abundant on simply-connected orbifolds, e.g. the double of a standard square.

By lifting to $(T_p, \tilde{g})$, we see that $\alpha * \beta$ is homotopic to $\alpha \cdot \beta$ via a *length decreasing* homotopy rel. end points. The assumption $|\alpha| + |\beta| < r_{\max}$ guarantees that the homotopy takes place in a domain of T_p on which $\exp_p$ is nonsingular. This is the orbifold version of the *short homotopy* in [BK].

Put the standard affine structure on $\mathbb{R}^n = T_p$. For each loop $\gamma \in \pi_\rho$ at p, we define the *holonomy motion*:

$$m(\gamma) : T_p \to T_p, \quad v \mapsto r(\gamma)v + t(\gamma). \tag{2.20}$$

Here $r(\gamma) \in O(n)$ is the parallel translation along γ, $t(\gamma) = |\gamma| \cdot \gamma'(|\gamma|)$. In another word, $m(\gamma)v = A(|\gamma|)$, where A is the solution of the following ODE along γ:

$$\begin{cases} \nabla_T A = T, \\ A(0) = v. \end{cases} \tag{2.21}$$

In particular, $m(\alpha \cdot \beta) = m(\alpha) \circ m(\beta)$. As in [BK], one can estimate the error caused by homotopy and conclude that

$$m(\alpha * \beta) = m(\alpha) \circ m(\beta) \quad \text{with error up to } \ c\Lambda^2\rho^2 < \epsilon_n \ll 1. \tag{2.22}$$

The following result of Gromov, [Gr], is of fundamental importance:

Proposition 2.23. *There is a positive constant* $\theta \ll 1$*, so that if* $|\alpha| \leq \rho/3$ *and* $|r(\alpha)| \leq 0.48$*, then*

$$\text{maximum rotational angle of } r(\alpha) \leq \theta\rho^{-1}|\alpha|. \tag{2.24}$$

In Chapter 3 of [BK], a detailed proof of the above proposition were given in the manifold case. All the arguments there (which involves Jacobi field estimates) were carried out on the tangent space T_p; therefore the proof in [BK] can be used here, by replacing the topological Gromov product in [BK] with the one we defined above.

Thus the rotation parts of $m(\gamma)$, with γ not too long, are concentrated in finitely many regions, $S_1, S_2, ..., S_N \subset O(n)$, which are mutually separated by distance at least 0.47. We can put these loops into finitely many classes, loops in the same class have rotational parts in the same S_i. There is a natural group structure on the set H of these classes, and H can be embedded into $O(n)$, with

$$|H| \leq (2\pi/0.47)^{\dim O(n)}. \tag{2.25}$$

Fix an orthonormal frame u at $0 \in (T_p, \tilde{g})$, parallel translate it along normal geodesics, we get an almost parallel frame field u on $B_\rho \subset T_p$. Let $q \in X$ be a regular point. Let $\tilde{q}_1, \tilde{q}_2 \in T_p$ so that $\exp(\tilde{q}_1) = \exp(\tilde{q}_2) = q$. Then there is an element $h \in H$, so that

$$d(\exp_*(u(\tilde{q}_1))h, \exp_*(u(\tilde{q}_2))) \leq \epsilon_n \ll 0.47. \tag{2.26}$$

In particular, if $\{\tilde{q}_1, \tilde{q}_2, ...\} = B_\rho(0) \cap \exp^{-1}(q)$, then the frames $\exp(u(\tilde{q}_i))$, $i = 1, 2, ...,$ also concentrated in regions $S_1^q, ..., S_N^q$. There is a right action by the discrete group H, which permutes these regions, with error up to $\epsilon_n \ll 0.47$. By a standard center of mass argument, for each $h \in H$ we get a frame $C_q(h)$ at q so that

$$C_q(h)h' = C_q(hh'), \qquad \text{for any } \ h' \in H. \tag{2.27}$$

Consider the set $X' \subset FX$ of all frames $C_q h$, denote by $\tilde{X}$ its closure. So $\operatorname{Diam} \tilde{X} \leq 2|H|$. Now it is easy to see that $\tilde{X}$ is a smooth manifold, and $X = H \backslash \tilde{X}$, see [BK]. Write $\pi : \tilde{X} \to X$ the projection. Any point $\tilde{q} \in \tilde{X}$ can also be viewed as a frame $v(\tilde{q})$ on X. Then $\pi^* v(\tilde{q})|_{\tilde{q}}$ gives a *global* orthonormal frame field on $\tilde{X}$.

Finally, following the deformation technique in Ruh, [Ru], and Ghanaat [Gh], we see $\tilde{X}$ is a nilmanifold, that concludes the proof of Gromov's theorem.

ACKNOWLEDGEMENTS. We are grateful to Prof. Tian for very helpful suggestions.

REFERENCES

[BK] Buser, P and Karcher, *Gromov's almost flat manifolds.* Astérisque 81, 1981.

[CG1] Cheeger, J; Gromov, M. Collapsing Riemannian manifolds while keeping their curvature bounded. I. J. Differential Geom. 23 (1986), no. 3, 309–346

[CG2] Cheeger, J; Gromov, M. *Collapsing Riemannian manifolds while keeping their curvature bounded.* II. J. Differential Geom. 32 (1990), no. 1, 269–298

[CT] Chen, X, Tian, G. *Ricci flow on Kähler-Einstein manifolds.* Duke Math. J. 131 (2006), no. 1, 17–73.

[CFG] Cheeger, J; Fukaya, K; Gromov, M. *Nilpotent structures and invariant metrics on collapsed manifolds*. J. Amer. Math. Soc. 5 (1992), no. 2, 327–372

[D] Ding, Y. *F-structure on collapsed orbifolds.* http://www.csulb.edu/~yding/orbifold.pdf

[F2] Fukaya, K. *A boundary of the set of the Riemannian manifolds with bounded curvatures and diameters*, J. Diff Geom. 28 (1988). 1-21.

[Gh] Ghanaat, P. *Almost Lie groups of type* $\mathbb{R}^n$. J. Reine Angew. Math. 401 (1989), 60–81.

[Gh2] Ghanaat, P. *Diskrete Gruppen und die Geometrie der Reprebndel*. J. Reine Angew. Math. 492 (1997), 135–178.

[Gr] Gromov, M. *Almost flat manifolds.* J. Diff Geom 13 (1978) 231-241.

[Ro] Rong, X. *Convergence and collapsing theorems in Riemannian geometry.* in Handbook of Geometric Analysis (Vol II), 193-299. International Press, 2010.

[Ru] Ruh, E. *Almost flat manifolds.* J. Diff Geom 17 (1982) 1-14.

[Th] Thurston, W. *The geometry and topology of* 3*-manifolds.* Princeton University. 1979.

Department of Mathematics and Statistics, California State University, Long Beach, 90840

E-mail address: `yding@csulb.edu`